arXiv:0805.0151v1 [math.AT] 1 May 2008

On the calculation of $H^*(\exp_3 S^n)$

S. C. F. Rose

May 1, 2008

Abstract

In this paper we provide two separate calculations for the cohomology of the space $\exp_3 S^n$. The first of these relies on a decomposition of this space into configuration spaces which have a large amount of geometric structure with which we can play, while the second proof will, with a bit of extra effort, provide a calculation of the cohomology of $\exp_k S^n$ for all k, n.

1 Introduction

The space of all non-empty finite subsets (of cardinality at most k) of a given space X, denoted at times $\exp_k(X), Sub_k(X), \mathcal{F}_k(X)$ or $X^{(k)}$, has been studied at varying frequencies since 1949 [Bor49]. The most celebrated result, of course, is in a correction to this initial paper wherein it is proven [Bot52] that $\exp_3 S^1 \cong S^3$. In a number of more recent papers [KS] [Tuf02] this result has been extended to show that $\exp_k(S^1)$ has the homotopy type of an odd dimensional sphere for all k.

In this paper we extend existing knowledge in the other direction—that is, we begin studying the spaces $\exp_3(S^n)$ for higher values of n.

For the duration of this paper all coefficients will be assumed to be $\mathbb{Z}/_2$, and the following notation will be used. $F_k(X)$ will refer to the orderd configuration space of k distinct points in X, while $C_k(X) := F_k(X)/_{\Sigma_k}$ will denote the unordered configuration space of points.

2 The Cohomology of $\exp_3(S^n)$

2.1 The Cohomology of $\exp_2(S^n)$

Lemma 2.1 *$C_2(S^n) \cong E$ where E is the orthogonal complement of the canonical line bundle over $\mathbb{R}P^n$.*

Proof Given a pair of points $\{x_1, x_2\}$ in S^n we find a line ℓ in $\mathbb{R}^{n+1}$. We then map those points to the line through the origin which is parallel to ℓ. The fibre of this map is then all pairs of points $\{x_1, x_2\}$ which are obtained from one another via a reflection in the plane perpendicular to ℓ. It is easy to see that this is the space E as claimed.



Figure 1: The map between $C_2(S^n)$ and E

Q.E.D.

It is worth noting that, being the orthogonal complement of the canonical line bundle, all of the Stiefel-Whitney classes of this bundles are non-zero.

To calculate the cohomology of $\exp_2(S^n)$, we begin with a Mayer-Vietoris sequence. Let U be the subset contained within $C_2(S^n)$ consisting of all points whose norm in the fibre is < 2, and let V be the subset of all those whose norm is > 1, together with $C_1(S^n) = S^n$. Then we have the following:

$$U \cong C_2(S^n) \simeq \mathbb{R}P^n$$
$$V \simeq S^n$$
$$U \cap V \simeq E_0$$

where E_0 is the sphere bundle contained in E.

We then have the long exact sequence

$$\cdots \to H^{k-1}(E_0) \to H^k(\exp_2 S^n) \to H^k(C_2 S^n) \oplus H^k(S^n) \to H^k(E_0) \to \cdots \tag{1}$$

which simplifies greatly to (as $H^k(C_2 S^n) \cong H^k(S^n) \cong 0$ for $k > n$)

$$H^k(E_0) \cong H^{k+1}(\exp_2 S^n)$$

for $k > n$. To study the remainder of this sequence we need to learn more about the cohomology of E_0.

Proposition 2.2 *With E_0 as above, we have*

$$H^*(E_0) \cong \langle \alpha, \beta \big| |\alpha| = 1, |\beta| = n, \alpha^n = 0 = \beta^2 \rangle$$

Proof Being a spherical fibration, with $\mathbb{Z}/_2$ coefficients we have the Gysin sequence

$$\cdots \to H^{k-1}(E_0) \to H^{k-n}(\mathbb{R}P^n) \xrightarrow{\smile e} H^k(\mathbb{R}P^n) \to H^k(E_0) \to \cdots$$

where the cup product is with the euler class of the fibration. But since the Euler class mod 2 is simply the top Stiefel-Whitney class—in this case, $\alpha^n \in H^n(\mathbb{R}P^n)$—the fact that all of our cohomology groups are $\mathbb{Z}/_2$ immediately follows. The product structure is an easy exercise after that.

Q.E.D.

Next, we note that in the long exact sequence of the pair (E, E_0) we find that the map $H^k(\mathbb{R}P^n) \to H^k(E_0)$ is by necessity an isomorphism for $k = 0, \ldots, n-2$ until we reach

$$0 \to H^{n-1}(\mathbb{R}P^n) \to H^{n-1}(E_0) \to H^0(\mathbb{R}P^n) \to \cdots$$

at which point it is an injection. However, this maps is the same as the map $H^k(C_2 S^n) \to H^k(E_0)$ in the Mayer-Vietoris sequence, and so we know that $H^k(\exp_2 S^n) = 0$ for $k = 1, \ldots, n-1$ leaving us with the exact sequence

$$0 \to H^n(\exp_2 S^n) \to H^n(C_2 S^n) \oplus H^n(S^n) \to H^n(E_0) \to H^{n+1}(\exp_2 S^n) \to 0$$

Now, for the map $H^n(C_2 S^n) \to H^n(E_0)$ we see from the Gysin sequence that it must be zero, since the sequence at that point looks like

$$\begin{tikzcd}
\cdots \arrow[r] & H^0(\mathbb{R}P^n) \arrow[r, "\smile \alpha^n"] \arrow[d, equal] & H^n(\mathbb{R}P^n) \arrow[r] \arrow[d, equal] & H^n(E_0) \arrow[r] \arrow[d, equal] & \cdots \\
& \mathbb{Z}/_2 \arrow[r, "1"] & \mathbb{Z}/_2 \arrow[r, "0"] & \mathbb{Z}/_2 &
\end{tikzcd}$$

As for the map $H^n(S^n) \to H^n(E_0)$, we can use the following geometric argument to show that it must be non-zero, and thus an isomorphism. We begin by providing an explicit description of the map $E_0 \to S^n$ which induces the map in the Mayer-Vietoris sequence.

As E_0 is the sphere bundle of the orthogonal complement to the canonical line bundle over $\mathbb{R}P^n$, we can consider a particular point $v \in E_0$ to be the corresponding point in S^n which lies on the circle perpendicular to $\pi(v)$, where π is the projection $E_0 \to \mathbb{R}P^n$. At this point the map $E_0 \to S^n$ is obvious—simply map a given point v to itself in S^n.

With this in mind, we can see that the "cell" consisting of $\pi^{-1}(\mathbb{R}P^1)$ maps via a degree one map onto S^n (where $\mathbb{R}P^1$ is the canonical $\mathbb{R}P^1 \hookrightarrow \mathbb{R}P^n$); as maps from this cell generate $H^n(E_0)$, it follows that the map $H^n(S^n) \to H^n(E_0)$ is non-trivial—in particular, it is an isomorphism—as claimed.

From all of the above we have

Proposition 2.3 *The mod 2 cohomology groups of* $\exp_2 S^n$ *are*

$$H^*(\exp_2 S^n) = \begin{cases} \mathbb{Z}/_2 & * = n, n+2, n+3, \ldots, 2n-1, 2n \\ 0 & \textit{otherwise} \end{cases}$$

Furthermore, the following can also be shown

Proposition 2.4 *If* e_i *(for* $i = n, n+2, n+3, \ldots, 2n$*) is the generator of* $H^i(\exp_2 S^n)$*, then we have that* $Sq^i(e_n) = e_{n+i}$ *for* $i = 2, \ldots, n$*.*

This is seen as a consequence of §3 of [Nak55] and the homological splitting of $H_n(Sym_k X)$.

2.2 The Main Calculation

We use a similar approach in this case as in the previous. Noting that any three points on a sphere S^n define an affine plane in $\mathbb{R}^{n+1}$, we find a map $C_3(S^n) \to \mathbb{G}_2\mathbb{R}^{n+1}$, the grassmannian of 2-planes in $\mathbb{R}^{n+1}$. This map is a fibration whose fibre contains, again, the orthogonal complement of the given plane. In this case though, the fibre is the direct sum of this orthogonal complement and a copy of $C_3(S^1)$ which we know is homeomorphic to a model Seifert fibration. Thus $C_3(S^n)$ is homotopic to a circle bundle over $\mathbb{G}_2\mathbb{R}^{n+1}$ and we may use the well known cohomology of this latter space, as well as a similar Mayer-Vietoris sequence to the one in the previous section to help determine the cohomology of $\exp_3(S^n)$.

The cohomology of $C_3(S^n)$ is relatively straightforward to determine. As a bundle over $\mathbb{G}_2\mathbb{R}^{n+1}$ with fibre homotopic to S^1, we have again the exact sequence

$$\cdots \to H^k(C_3 S^n) \to H^{k-1}(\mathbb{G}_2\mathbb{R}^{n+1}) \xrightarrow{\smile\alpha} H^{k+1}(\mathbb{G}_2\mathbb{R}^{n+1}) \to H^{k+1}(C_3 S^n) \to \cdots \tag{2}$$

Now, as in [Bor53], we know that

$$H^*(\mathbb{G}_2\mathbb{R}^{n+1}) \cong \mathbb{Z}/_2[\alpha_1, \alpha_2, \beta_1, \ldots, \beta_{n-1}]/\langle (1 + \sum \alpha_i)(1 + \sum \beta_i) = 1 \rangle \tag{3}$$

with $\alpha_i, \beta_i \in H^i(\mathbb{G}_2\mathbb{R}^{n+1})$ being the Stiefel-Whitney classes of the canonical 2-plane bundle and dual $(n-1)$-plane bundles, respectively.

We first claim that $C_3 S^n$ is homotopic to the circle bundle of the canonical 2-plane bundle over $\mathbb{G}_2\mathbb{R}^{n+1}$—as such, the α in the exact sequence (2) is nothing but α_2. Noting that the fibre of this bundle, $\mathbb{R}^{n-1} \times C_3 S^1$ deformation retracts onto the central fibre of $C_3 S^1$ we obtain the first part. But this central fibre consists of triples of points which form an equilateral triangle in S^1. It is easy to see that such a bundle is simply equivalent to an ordinary S^1 bundle by taking the fibre-wise map $z \mapsto z^3$. The deformation in question also translates our plane to the origin, at which point the conclusion follows.

We can in fact say a little more.

Proposition 2.5 *The space C_3S^n, as a bundle over $\mathbb{G}_2\mathbb{R}^{n+1}$, is the fibrewise direct product of γ_2, γ_{n-1} and S^1, where the γ_i are the tautological and dual bundles over $\mathbb{G}_2\mathbb{R}^{n+1}$, respectively, and S^1 is the sphere bundle of γ_2.*

Proof As above, we note that this is in fact the fibrewise direct product of γ_{n-1} and C_3S^1. Now, C_3S^1 is a quotient of $F_3S^1 \cong B^2 \times S^1$ by the action of the cyclic group $\mathbb{Z}/_3$ (see [Ros07]). From all of this we can consider C_3S^n to be the quotient of the bundle $\gamma_{n-1} \oplus (B^2 \times S^1)$ by a $\mathbb{Z}/_3$ action.

Now, since the S^1 component is, as above, the sphere bundle of the tautological 2-plane bundle over $\mathbb{G}_2\mathbb{R}^{n+1}$, we can write out an explicit homeomorphism between this space and $\gamma_{n-1} \oplus \gamma_2 \oplus S^1$. However, on the fibres (see again [Ros07]) we see that the action of $\mathbb{Z}/_3$ yields that this space is simply $\gamma_{n-1} \oplus \left(\gamma_2/\mathbb{Z}/_3\right) \oplus S^1$, from which the conclusion follows.

Q.E.D.

Corollary 2.6 *$C_3S^n \cong \varepsilon^{n+1} \oplus S^1$, where S^1 is the sphere bundle of γ_2, the tautological bundle over $\mathbb{G}_2\mathbb{R}^{n+1}$.*

To understand the resulting cohomology of C_3S^n, we examine that of $\mathbb{G}_2\mathbb{R}^{n+1}$ further. The first thing to note is that the relation(s) given in (3) allow us to write all of the generators β_i in terms of α_1, α_2. For example, we note that we can write the first few terms as

$$\begin{aligned} \alpha_1 + \beta_1 &= 0 \\ \alpha_2 + \alpha_1\beta_1 + \beta_2 &= 0 \\ \alpha_2\beta_1 + \alpha_1\beta_2 + \beta_3 &= 0 \end{aligned}$$

from which it follows that $\beta_1 = \alpha_1, \beta_2 = \alpha_2 + \alpha_1^2$ and $\beta_3 = \alpha_1^3$. Now, the first relation which isn't simply the elimination of extraneous generators is in dimension n (In particular, it is the relation that $\beta_{n-1}\alpha_1 + \beta_{n-2}\alpha_2 = 0$). As such, up until dimension n, the dimension of our cohomology groups is simply the number of unique products of the α_1, α_2. To wit,

i	0	1	2	3	4	5	6	$\cdots$
dim $H^i(\mathbb{G}_2\mathbb{R}^{n+1})$	1	1	2	2	3	3	4	$\cdots$

As $\mathbb{G}_2\mathbb{R}^{n+1}$ is a manifold of dimension $2n-2$ (and, more importantly, we are working with $\mathbb{Z}/_2$ coefficients), Poincaré duality holds and we see that $H^{2n-2-i}(\mathbb{G}_2\mathbb{R}^{n+1}) \cong H^i(\mathbb{G}_2\mathbb{R}^{n+1})$.

Moroever, we can show that up until (and including) dimension $k = n-1$ the map $H^{k-2}(\mathbb{G}_2\mathbb{R}^{n+1}) \xrightarrow{\smile\alpha_2} H^k(\mathbb{G}_2\mathbb{R}^{n+1})$ is an injection (there are no relations up until that dimension). We also have that $H^{n-2}(\mathbb{G}_2\mathbb{R}^{n+1}) \xrightarrow{\smile\alpha_2} H^n(\mathbb{G}_2\mathbb{R}^{n+1})$ is an isomorphism, and that beyond that it is a surjective map. We then have the following.

$$\begin{tikzcd}
n+1 & 1 \arrow[drr, "\smile\alpha_2"'] & \langle\alpha_1\rangle \arrow[drr, "\smile\alpha_2"'] & \langle\alpha_1^2,\alpha_2\rangle \arrow[drr, "\smile\alpha_2"'] & \langle\alpha_1^3,\alpha_2\alpha_1\rangle \arrow[drr, "\smile\alpha_2"'] & \langle\alpha_1^4,\alpha_2\alpha_1^2,\alpha_2^2\rangle & \cdots \\
n & 1 & \langle\alpha_1\rangle & \langle\alpha_1^2,\alpha_2\rangle & \langle\alpha_1^3,\alpha_2\alpha_1\rangle & \langle\alpha_1^4,\alpha_2\alpha_1^2,\alpha_2^2\rangle & \cdots \\
\vdots & & & & & & \\
1 & 1 \arrow[drr, "\smile\alpha_2"'] & \langle\alpha_1\rangle \arrow[drr, "\smile\alpha_2"'] & \langle\alpha_1^2,\alpha_2\rangle \arrow[drr, "\smile\alpha_2"'] & \langle\alpha_1^3,\alpha_2\alpha_1\rangle \arrow[drr, "\smile\alpha_2"'] & \langle\alpha_1^4,\alpha_2\alpha_1^2,\alpha_2^2\rangle & \cdots \\
0 & 1 & \langle\alpha_1\rangle & \langle\alpha_1^2,\alpha_2\rangle & \langle\alpha_1^3,\alpha_2\alpha_1\rangle & \langle\alpha_1^4,\alpha_2\alpha_1^2,\alpha_2^2\rangle & \cdots
\end{tikzcd}$$

Figure 2: $E_2^{*,*}$

Proposition 2.7 *The mod 2 cohomology ring of $C_3(S^n)$, the collection of unorderd, distinct triples of points in S^n is given by*

$$H^*(C_3S^n) \cong \langle \alpha, \beta | \alpha^n = \beta^2 = 0, |\alpha| = 1, |\beta| = n \rangle$$

We next examine the more complicated portion—the intersection of the two parts, which is a bundle over $\mathbb{G}_2\mathbb{R}^{n+1}$ with fibre $F := \partial(B^{n-1} \times C_3S^1)$.

To compute the cohomology of this part, we use the Leray-Serre spectral sequence. The first bit of note is that the cohomology of the fibre is given by

$$H^*(F) = \langle \alpha, \beta | \alpha^2 = \beta^2 = 0, |\alpha| = 1, |\beta| = n \rangle$$

With this in mind, the E_2 term of the spectral sequence is shown in figure 2

Moreover, by the same reasoning used in the construction of the Gysin sequence, we see that $d_2(\alpha)$ is simply the cup product with the euler class of the spherical fibration represented by the class α—and thus that it is simply the cup product with α_2, as in the calculation for C_3S^n. If $n \neq 2$, then $d_2(\beta) = 0$, and so we have the E_n sheet looking like the one in figure 3

The only possible non-zero differential of note is $d_n(\beta)$, which we claim must be zero. That being the case, we have the following.

Proposition 2.8 *The mod 2 cohomology of the total space of the bundle $\partial(B^{n-1} \times C_3S^1) \hookrightarrow E \to \mathbb{G}_2\mathbb{R}^{n+1}$ is given by*

$$H^*(E) = \begin{cases} \mathbb{Z}/_2 & * = 0, \ldots, n-1, 2n, \ldots, 3n-1 \\ \mathbb{Z}/_2 \oplus \mathbb{Z}/_2 & * = n, \ldots, 2n-1 \\ 0 & \textit{otherwise} \end{cases}$$



Figure 3: $E_n^{*,*}$

Proof of claim: The proof is quite simple. From corollary 2.6, we know that $C_3S^n \cong \varepsilon^{n+1} \oplus S^1$. As a consequence, we have a projection from this bundle to the S^1 factor. Since this bundle is a trivial bundle with an addition S^1 part, we also have a section of this bundle. Thus simple algebraic arguments yeild that there must be non-zero cohomology classes in $H^{n+1}(E)$, which would be zero if $d_n(\beta)$ were non-zero.

Q.E.D.

Corollary 2.9 *The map $H^*(C_3S^n) \to H^*(E)$ is injective.*

We are now ready to put this all together to obtain the cohomology of $\exp_3(S^n)$. Recall that we have the Mayer-Vietoris sequence

$$\begin{aligned}\cdots \to H^k(\exp_3 S^n) \to H^k(C_3S^n) \oplus H^k(\exp_2 S^n) \to \\ \to H^k(E) \to H^{k+1}(\exp_3 S^n) \to \cdots \quad (4)\end{aligned}$$

where the cohomology of each known term is described in propositions 2.7, 2.3 and 2.8, respectively. If we consider corollary 2.9, then all that remains is determining the map $H^*(\exp_2 S^n) \to H^*(E)$.

In dimension n this is an isomorphism. From [KS] we note that $\exp_3(X)$ is $(r+1)$-connected where X is r-connected. In our case, this means that $\exp_3(S^n)$ is n-connected, and so its n-th homology group is necessarily trivial—but the Mayer-vietoris sequence

$$0 \to H^n(\exp_3 S^n) \to H^n(C_3S^n) \oplus H^n(\exp_2 S^n) \to H^n(E) \to H^{n+1}(\exp_3 S^n) \to 0$$

also yields that $H^{n+1}(\exp_3 S^n)$ is also zero.

What about higher dimensions? We note that by necessity $H^{n+2}(\exp_3 S^n)$ has at least one $\mathbb{Z}/_2$ factor in it, due to the sequence

$$\begin{tikzcd}
0 \arrow[r] & H^{n+1}(C_3S^n) \arrow[r, hook] \arrow[d, equal] & H^{n+1}(E) \arrow[r] \arrow[d, equal] & H^{n+2}(\exp_3 S^n) \\
& \mathbb{Z}/_2 \arrow[r, hook] & \mathbb{Z}/_2 \oplus \mathbb{Z}/_2 \arrow[ur] &
\end{tikzcd}$$

Claim: The map $H^{n+k}(\exp_2 S^n) \to H^{n+k}(E)$ for $2 \le k \le n$ is 0. With this in mind, we can read off all of our cohomology groups of $\exp_3 S^n$ as follows.

Theorem 2.10 *The mod 2 cohomology groups of* $\exp_3 S^n$*, the collection of non-empty subsets of* S^n *of cardinality at most 3, is given by*

$$H^*(\exp_3 S^n) = \begin{cases} \mathbb{Z}/_2 & * = 0, 2n+1, 2n+2, \dots, 3n \\ \mathbb{Z}/_2 \oplus \mathbb{Z}/_2 & * = n+2, \dots, 2n \\ 0 & \textit{otherwise} \end{cases}$$

In particular, the cohomology of $\exp_3 S^2$ *is given by*

$$H^*(\exp_3 S^n) = \begin{cases} \mathbb{Z}/_2 \oplus \mathbb{Z}/_2 & * = 4 \\ \mathbb{Z}/_2 & * = 0, 5, 6 \\ 0 & \textit{otherwise} \end{cases}$$

Proof of claim: To prove this claim it will suffice to show that the map $Sq^i : H^n(E) \to H^{n+i}(E)$ is 0 for $2 \le i \le n$, since we have the diagram

$$\begin{tikzcd}
H^n(\exp_2 S^n) \arrow[r, hook] \arrow[d, "Sq^i", "\cong"'] & H^n(E) \arrow[d, "Sq^i"] \\
H^{n+i}(\exp_2 S^n) \arrow[r] & H^{n+i}(E)
\end{tikzcd}$$

Now, from Proposition 2.4 we know that Sq^i is an isomorphism $H^n(\exp_2 S^n) \to H^{n+i}(\exp_2 S^n)$ for $2 \le i \le n$. Since the map $H^n(C_3S^n) \oplus H^n(\exp_2 S^n) \to H^n(E)$ is an isomorphism in dimension n, it follows that all of the maps $H^{n+i}(C_3S^n) \oplus H^{n+i}(\exp_2 S^n) \to H^{n+i}(E)$ are determined by the Squares' actions on $\beta \in H^n(E)$—but this action is trivial for $2 \le i \le n$, since this action arises from the operators (See [McC01]) ${}_ASq^i : E_*^{p,q} \to E_*^{p,q+i}$ and ${}_BSq^i : E_*^{p,q} \to E_*^{p-q+i,2q}$. It is clear that, on any class in $E_n^{0,n}$, these are 0. As these converge to the action of $\mathcal{A}_2$, we see that $Sq^i(\beta) = 0$ as claimed.

Q.E.D.

3 An Alternate Proof

There is another way in which we can compute this cohomology which, while it offers less geometric insight, it does have the added benefit that it provides (in principle!) the first step in an inductive computation of $H^*(\exp_k S^n)$ for any k, n.

The ingredients for this proof are as follows (See [KS], [Nak55]). We have the pushout square

$$\begin{tikzcd}
S^n \times S^n \arrow[r, "\alpha"] \arrow[d, "q"] & Sym_3 S^n \arrow[d] \\
\exp_2 S^n \arrow[r, hook] & \exp_3 S^n
\end{tikzcd} \tag{5}$$

where the map α takes $(x, y) \mapsto x^2 y$, and q is the obvious quotient map. We also, as per Proposition 2.4 and [Nak55] have that

$$Sq^n(e_n) = e_{2n} \qquad Sq^n(f_n) = f_{2n}$$

where e_i, f_i denote the generators of $H^i(\exp_2 S^n)$ and $H^i(Sym_3 S^n)$, respectively.

Now, we consider the Mayer-Vietoris sequence which arises from the pushout (5), which reads as follows.

$$\cdots \to H^k(\exp_3 S^n) \to H^k(\exp_2 S^n) \oplus H^k(Sym_3 S^n) \to H^k(S^n \times S^n) \to \cdots$$

Fortunately, the last space in this row has mostly zero cohomology, and so a lot of isomorphisms $H^k(\exp_3 S^n) \cong H^k(Sym_3 S^n) \oplus H^k(\exp_2 S^n)$ immediately fall out. These only fail around dimensions $n, 2n$ where we have to be a little more careful.

These two sequences are as follows.

$$0 \to H^n(\exp_3 S^n) \to \mathbb{Z}/_2 \oplus \mathbb{Z}/_2 \to \mathbb{Z}/_2 \oplus \mathbb{Z}/_2 \to H^{n+1}(\exp_3 S^n) \to 0$$

and

$$\begin{aligned} 0 \to H^{2n}(\exp_3 S^n) \to H^{2n}(\exp_2 S^n) \oplus H^{2n}(Sym_3 S^n) \to \\ \to H^{2n}(S^n \times S^n) \to H^{2n+1}(\exp_3 S^n) \to H^{2n+1}(Sym_3 S^n) \to 0 \end{aligned}$$

The same connectivity argument as before shows that $H^n(\exp_3 S^n)$ (and hence $H^{n+1}(\exp_3 S^n)$ is zero—and moreover that the maps $H^n(\exp_2 S^n) \to H^n(S^n \times S^n)$ and $H^n(Sym_3 S^n) \to H^n(S^n \times S^n)$ are both injections.

Now, we look at the commuting squares

$$\begin{tikzcd}
H^n(\exp_2 S^n) \arrow[r] \arrow[d, "Sq^n", "\cong"'] & H^n(S^n \times S^n) \arrow[d, "Sq^n"] \\
H^{2n}(\exp_2 S^n) \arrow[r] & H^{2n}(S^n \times S^n)
\end{tikzcd}$$

and

$$\begin{tikzcd}
H^n(Sym_3 S^n) \arrow[r] \arrow[d, "\cong"', "Sq^n"] & H^n(S^n \times S^n) \arrow[d, "Sq^n"] \\
H^{2n}(Sym_3 S^n) \arrow[r] & H^{2n}(S^n \times S^n)
\end{tikzcd}$$

Since the squares act trivially on $H^*(S^n \times S^n)$ we see that the maps in dimension $2n$ must also be zero. The conclusion then follows.

To continue this argument, we note that we always have an analagous pushout to (5), namely

$$\begin{tikzcd}
(S^n)^{\times k} \arrow[r, "\alpha"] \arrow[d, "q"] & Sym_{k+1} S^n \arrow[d] \\
\exp_k S^n \arrow[r, hook] & \exp_{k+1} S^n
\end{tikzcd}$$

Thus all that remains is knowing the cohomology of the higher symmetric products of S^n, as well as determining the action of the Steenrod algebra on these spaces and on $\exp_k S^n$.

References

[Bor49] Karol Borsuk. On the third symmetric potency of the circumference. *Fund. Math.*, 36:236–244, 1949.

[Bor53] A. Borel. La cohomologie mod 2 de certains espaces homogènes. *Comment. Math. Helv.*, 27:165–197, 1953.

[Bot52] R. Bott. On the third symmetric potency of S_1. *Fund. Math.*, 39:264–268 (1953), 1952.

[KS] Sadok Kallel and Denis Sjerve. Finite subset spaces and a spectral sequence of Biro. preprint.

[McC01] John McCleary. *A User's guide to Spectral sequences.* Cambridge studies in advanced mathematics, 2001.

[Nak55] Minoru Nakaoka. Cohomology of the three-fold symmetric products of spheres. *Proc. Japan. Acad.*, 31:670–672, 1955.

[Ros07] Simon Rose. Finite subsets of the circle—a hypberbolic approach. Master's thesis, University of British Columbia, 2007.

[Tuf02] Christopher Tuffley. Finite subset spaces of S^1. *Algebr. Geom. Topol.*, 2:1119–1145 (electronic), 2002.

[Tuf03] Christopher Tuffley. Finite subset spaces of closed surfaces, 2003.